Overview of State Reporting of the 2025 Adult Core Set Measures

	Number of Core Set Measures Reported	Number of Provisional Measures Reported	Initiation and Engagement of Substance Use Disorder Treatment (IET-AD)*	Use of Pharmacotherapy for Opioid Use Disorder (OUD-AD)*	Medical Assistance with Smoking and Tobacco Use Cessation (MSC-AD)*	Follow-Up After ED Visit for Substance Use: Age 18 and Older (FUA-AD)*	Follow-Up After ED Visit for Mental Illness: Age 18 and Older (FUM-AD)*	Follow-Up After Hospitalization for Mental Illness: Age 18 and Older (FUH-AD)*	Antidepressant Medication Management (AMM-AD)*	Adherence to Antipsychotic Medications for Individuals With Schizophrenia (SAA-AD)*	Diabetes Screening for People With Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medications (SSD-AD)*	Diabetes Care for People with Serious Mental Illness: Glycemic Status (>9.0%) (HPCMI-AD)*	Screening for Depression and Follow-Up Plan: Age 18 and Older (CDF-AD)*	Breast Cancer Screening (BCS-AD)	Cervical Cancer Screening (CCS-AD)	Colorectal Cancer Screening (COL-AD)	Chlamydia Screening in Women Ages 21 to 24 (CHL-AD)	Adult Immunization Status (AIS-AD)	Prenatal and Postpartum Care: Age 21 and Older (PPC2-AD)	Prenatal Immunization Status: Age 21 and Older (PRS-AD)	Low-Risk Cesarean Delivery: Age 20 and Older (LRCD-AD)	Contraceptive Care – Postpartum Women Ages 21 to 44 (CCP-AD)	Contraceptive Care – All Women Ages 21 to 44 (CCW-AD)	Glycemic Status Assessment for Patients with Diabetes (GSD-AD)	PQI 01: Diabetes Short-Term Complications Admission Rate (PQI01-AD)	PQI 05: Chronic Obstructive Pulmonary Disease (COPD) or Asthma in Older Adults Admission Rate (PQI05-AD)	PQI 08: Heart Failure Admission Rate (PQI08-AD)	PQI 15: Asthma in Younger Adults Admission Rate (PQI15-AD)	Plan All-Cause Readmissions (PCR-AD)	Avoidance of Antibiotic Treatment for Acute Bronchitis/Bronchiolitis: Age 18 and Older (AAB-AD)	Asthma Medication Ratio: Ages 19 to 64 (AMR-AD)	Controlling High Blood Pressure (CBP-AD)	HIV Viral Load Suppression (HVL-AD)	Use of Opioids at High Dosage in Persons Without Cancer (OHD-AD)	Concurrent Use of Opioids and Benzodiazepines (COB-AD)	Oral Evaluation During Pregnancy: Ages 21 to 44 (OEVP-AD)	Ambulatory Care Sensitive ED Visits for Non-Traumatic Dental Conditions in Adults (EDV-AD)	CAHPS Health Plan Survey 5.1H, Adult Version (Medicaid) (CPA-AD)	Long-Term Services and Supports Comprehensive Care Plan and Update (CPU-AD)	National Core Indicators Survey (NCIIDD-AD)	Postpartum Depression Screening and Follow-Up: Age 21 and Older (PDS-AD)**
Total			**52**	**52**	**51**	**52**	**52**	**52**	**52**	**52**	**52**	**49**	**49**	**44**	**45**	**45**	**45**	**34**	**44**	**30**	**54**	**45**	**45**	**40**	**41**	**40**	**40**	**40**	**41**	**46**	**44**	**43**	**21**	**39**	**41**	**27**	**21**	**50**	**5**	**39**	**16**
Alabama	36	0	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	X	X	X	X	X	-	X	-
Alaska	13	0	X	X	X	X	X	X	X	X	X	X	X	-	-	-	-	-	-	-	X	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	X	-	-	-
Arizona	36	1	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	X	X	X	-	X	X	X	X
Arkansas	30	0	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	-	-	X	X	X	X	X	X	X	X	X	X	-	X	-	X	X	-	-	X	-	X	-
California	37	0	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	X	-
Colorado	36	1	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	X	-	X	X
Connecticut	35	0	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	X	X	X	-	X	-	X	-
Delaware	35	1	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	-	X	X	X	X
Dist. Of Col.	28	1	X	X	X	X	X	X	X	X	X	-	X	X	X	X	X	X	X	X	X	X	X	X	-	-	-	-	-	X	X	X	-	X	X	-	-	X	-	X	X
Florida	27	0	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	X	-	X	X	X	X	-	-	-	-	X	X	X	X	-	X	X	-	-	X	-	-	-
Georgia	14	0	X	X	X	X	X	X	X	X	X	X	X	-	-	-	-	-	-	-	X	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	X	-	X	-
Guam	1	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	X	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Hawaii	33	1	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	-	X	X	X	X	X	X	X	X	X	X	X	X	X	-	X	X	-	-	X	X	X	X
Idaho	32	0	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	X	-	X	X	X	X	X	X	X	X	-	X	X	X	-	X	X	X	X	X	-	-	-
Illinois	14	0	X	X	X	X	X	X	X	X	X	X	X	-	-	-	-	-	-	-	X	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	X	-	X	-

Overview of State Reporting of the 2025 Adult Core Set Measures (*continued*)

	Number of Core Set Measures Reported	Number of Provisional Measures Reported	Initiation and Engagement of Substance Use Disorder Treatment (IET-AD)*	Use of Pharmacotherapy for Opioid Use Disorder (OUD-AD)*	Medical Assistance with Smoking and Tobacco Use Cessation (MSC-AD)*	Follow-Up After ED Visit for Substance Use: Age 18 and Older (FUA-AD)*	Follow-Up After ED Visit for Mental Illness: Age 18 and Older (FUM-AD)*	Follow-Up After Hospitalization for Mental Illness: Age 18 and Older (FUH-AD)*	Antidepressant Medication Management (AMM-AD)*	Adherence to Antipsychotic Medications for Individuals With Schizophrenia (SAA-AD)*	Diabetes Screening for People With Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medications (SSD-AD)*	Diabetes Care for People with Serious Mental Illness: Glycemic Status (>9.0%) (HPCMI-AD)*	Screening for Depression and Follow-Up Plan: Age 18 and Older (CDF-AD)*	Breast Cancer Screening (BCS-AD)	Cervical Cancer Screening (CCS-AD)	Colorectal Cancer Screening (COL-AD)	Chlamydia Screening in Women Ages 21 to 24 (CHL-AD)	Adult Immunization Status (AIS-AD)	Prenatal and Postpartum Care: Age 21 and Older (PPC2-AD)	Prenatal Immunization Status: Age 21 and Older (PRS-AD)	Low-Risk Cesarean Delivery: Age 20 and Older (LRCD-AD)	Contraceptive Care – Postpartum Women Ages 21 to 44 (CCP-AD)	Contraceptive Care – All Women Ages 21 to 44 (CCW-AD)	Glycemic Status Assessment for Patients with Diabetes (GSD-AD)	PQI 01: Diabetes Short-Term Complications Admission Rate (PQI01-AD)	PQI 05: Chronic Obstructive Pulmonary Disease (COPD) or Asthma in Older Adults Admission Rate (PQI05-AD)	PQI 08: Heart Failure Admission Rate (PQI08-AD)	PQI 15: Asthma in Younger Adults Admission Rate (PQI15-AD)	Plan All-Cause Readmissions (PCR-AD)	Avoidance of Antibiotic Treatment for Acute Bronchitis/Bronchiolitis: Age 18 and Older (AAB-AD)	Asthma Medication Ratio: Ages 19 to 64 (AMR-AD)	Controlling High Blood Pressure (CBP-AD)	HIV Viral Load Suppression (HVL-AD)	Use of Opioids at High Dosage in Persons Without Cancer (OHD-AD)	Concurrent Use of Opioids and Benzodiazepines (COB-AD)	Oral Evaluation During Pregnancy: Ages 21 to 44 (OEVP-AD)	Ambulatory Care Sensitive ED Visits for Non-Traumatic Dental Conditions in Adults (EDV-AD)	CAHPS Health Plan Survey 5.1H, Adult Version (Medicaid) (CPA-AD)	Long-Term Services and Supports Comprehensive Care Plan and Update (CPU-AD)	National Core Indicators Survey (NCIIDD-AD)	Postpartum Depression Screening and Follow-Up: Age 21 and Older (PDS-AD)**
Indiana	31	0	X	X	X	X	X	X	X	X	X	X	-	X	X	X	X	X	X	-	X	X	X	X	X	X	X	X	X	X	X	X	-	-	X	-	-	X	-	X	-
Iowa	32	0	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	X	-	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	-	X	-	-	-
Kansas	38	0	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-
Kentucky	30	0	X	X	X	X	X	X	X	X	X	X	X	-	X	-	X	-	X	-	X	X	X	-	X	X	X	X	X	X	X	X	X	X	X	-	-	X	-	X	-
Louisiana	34	0	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	X	-	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	X	-	X	-
Maine	32	0	X	X	-	X	X	X	X	X	X	-	-	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	-	-	-
Maryland	35	0	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	X	-	X	-
Massachusetts	34	1	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	X	X	X	-	X	-	-	X
Michigan	36	0	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	X	-	X	-
Minnesota	35	0	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	X	X	X	X	-	-	X	-
Mississippi	35	0	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	X	X	X	X	X	-	-	-
Missouri	35	0	X	X	X	X	X	X	X	X	X	-	-	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	X	-
Montana	19	1	X	X	X	X	X	X	X	X	X	X	X	X	-	-	-	-	-	-	X	-	-	-	-	-	-	-	-	-	-	X	-	X	X	-	X	X	-	X	X
Nebraska	37	0	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	X	-
Nevada	23	0	X	X	X	X	X	X	X	X	X	X	X	-	-	X	-	-	-	-	X	X	X	-	X	X	X	X	-	X	-	-	X	-	-	-	-	X	-	X	-

Overview of State Reporting of the 2025 Adult Core Set Measures (*continued*)

	Number of Core Set Measures Reported	Number of Provisional Measures Reported	Initiation and Engagement of Substance Use Disorder Treatment (IET-AD)*	Use of Pharmacotherapy for Opioid Use Disorder (OUD-AD)*	Medical Assistance with Smoking and Tobacco Use Cessation (MSC-AD)*	Follow-Up After ED Visit for Substance Use: Age 18 and Older (FUA-AD)*	Follow-Up After ED Visit for Mental Illness: Age 18 and Older (FUM-AD)*	Follow-Up After Hospitalization for Mental Illness: Age 18 and Older (FUH-AD)*	Antidepressant Medication Management (AMM-AD)*	Adherence to Antipsychotic Medications for Individuals With Schizophrenia (SAA-AD)*	Diabetes Screening for People With Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medications (SSD-AD)*	Diabetes Care for People with Serious Mental Illness: Glycemic Status (>9.0%) (HPCMI-AD)*	Screening for Depression and Follow-Up Plan: Age 18 and Older (CDF-AD)*	Breast Cancer Screening (BCS-AD)	Cervical Cancer Screening (CCS-AD)	Colorectal Cancer Screening (COL-AD)	Chlamydia Screening in Women Ages 21 to 24 (CHL-AD)	Adult Immunization Status (AIS-AD)	Prenatal and Postpartum Care: Age 21 and Older (PPC2-AD)	Prenatal Immunization Status: Age 21 and Older (PRS-AD)	Low-Risk Cesarean Delivery: Age 20 and Older (LRCD-AD)	Contraceptive Care – Postpartum Women Ages 21 to 44 (CCP-AD)	Contraceptive Care – All Women Ages 21 to 44 (CCW-AD)	Glycemic Status Assessment for Patients with Diabetes (GSD-AD)	PQI 01: Diabetes Short-Term Complications Admission Rate (PQI01-AD)	PQI 05: Chronic Obstructive Pulmonary Disease (COPD) or Asthma in Older Adults Admission Rate (PQI05-AD)	PQI 08: Heart Failure Admission Rate (PQI08-AD)	PQI 15: Asthma in Younger Adults Admission Rate (PQI15-AD)	Plan All-Cause Readmissions (PCR-AD)	Avoidance of Antibiotic Treatment for Acute Bronchitis/Bronchiolitis: Age 18 and Older (AAB-AD)	Asthma Medication Ratio: Ages 19 to 64 (AMR-AD)	Controlling High Blood Pressure (CBP-AD)	HIV Viral Load Suppression (HVL-AD)	Use of Opioids at High Dosage in Persons Without Cancer (OHD-AD)	Concurrent Use of Opioids and Benzodiazepines (COB-AD)	Oral Evaluation During Pregnancy: Ages 21 to 44 (OEVP-AD)	Ambulatory Care Sensitive ED Visits for Non-Traumatic Dental Conditions in Adults (EDV-AD)	CAHPS Health Plan Survey 5.1H, Adult Version (Medicaid) (CPA-AD)	Long-Term Services and Supports Comprehensive Care Plan and Update (CPU-AD)	National Core Indicators Survey (NCIIDD-AD)	Postpartum Depression Screening and Follow-Up: Age 21 and Older (PDS-AD)**
New Hampshire	36	1	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	X	X	X	X	X	-	X	X
New Jersey	37	1	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	X	X
New Mexico	28	1	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	-	-	X	X	X	X	-	-	-	-	-	X	-	-	X
New York	34	0	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	-	X	-	X	-
North Carolina	36	0	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	X	X	X	X	X	-	X	-
North Dakota	37	1	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	X	X
Ohio	30	0	X	X	X	X	X	X	X	X	X	X	X	X	X	-	X	-	X	-	X	X	X	X	X	X	X	X	X	X	X	X	-	X	-	-	-	X	-	X	-
Oklahoma	34	0	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	-	X	X	-	X	X	-	X	-
Oregon	27	0	X	X	X	X	X	X	X	X	X	X	X	-	X	X	X	-	X	-	X	X	X	-	-	-	-	-	X	X	X	-	-	X	X	X	X	X	-	X	-
Pennsylvania	36	1	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	X	X	X	X	X	-	X	X
Puerto Rico	36	1	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	-	X
Rhode Island	25	1	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	-	X	-	-	-	-	X	X	X	X	-	-	-	-	-	X	-	-	X
South Carolina	36	0	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	X	X	X	X	X	-	X	-
South Dakota	24	0	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	X	X	X	-	-	-	-	-	-	X	X	-	-	-	-	-	-	X	-	X	-
Tennessee	36	1	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	X	X	X	X	X	X	-	X

Overview of State Reporting of the 2025 Adult Core Set Measures (*continued*)

	Number of Core Set Measures Reported	Number of Provisional Measures Reported	Initiation and Engagement of Substance Use Disorder Treatment (IET-AD)*	Use of Pharmacotherapy for Opioid Use Disorder (OUD-AD)*	Medical Assistance with Smoking and Tobacco Use Cessation (MSC-AD)*	Follow-Up After ED Visit for Substance Use: Age 18 and Older (FUA-AD)*	Follow-Up After ED Visit for Mental Illness: Age 18 and Older (FUM-AD)*	Follow-Up After Hospitalization for Mental Illness: Age 18 and Older (FUH-AD)*	Antidepressant Medication Management (AMM-AD)*	Adherence to Antipsychotic Medications for Individuals With Schizophrenia (SAA-AD)*	Diabetes Screening for People With Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medications (SSD-AD)*	Diabetes Care for People with Serious Mental Illness: Glycemic Status (>9.0%) (HPCMI-AD)*	Screening for Depression and Follow-Up Plan: Age 18 and Older (CDF-AD)*	Breast Cancer Screening (BCS-AD)	Cervical Cancer Screening (CCS-AD)	Colorectal Cancer Screening (COL-AD)	Chlamydia Screening in Women Ages 21 to 24 (CHL-AD)	Adult Immunization Status (AIS-AD)	Prenatal and Postpartum Care: Age 21 and Older (PPC2-AD)	Prenatal Immunization Status: Age 21 and Older (PRS-AD)	Low-Risk Cesarean Delivery: Age 20 and Older (LRCD-AD)	Contraceptive Care – Postpartum Women Ages 21 to 44 (CCP-AD)	Contraceptive Care – All Women Ages 21 to 44 (CCW-AD)	Glycemic Status Assessment for Patients with Diabetes (GSD-AD)	PQI 01: Diabetes Short-Term Complications Admission Rate (PQI01-AD)	PQI 05: Chronic Obstructive Pulmonary Disease (COPD) or Asthma in Older Adults Admission Rate (PQI05-AD)	PQI 08: Heart Failure Admission Rate (PQI08-AD)	PQI 15: Asthma in Younger Adults Admission Rate (PQI15-AD)	Plan All-Cause Readmissions (PCR-AD)	Avoidance of Antibiotic Treatment for Acute Bronchitis/Bronchiolitis: Age 18 and Older (AAB-AD)	Asthma Medication Ratio: Ages 19 to 64 (AMR-AD)	Controlling High Blood Pressure (CBP-AD)	HIV Viral Load Suppression (HVL-AD)	Use of Opioids at High Dosage in Persons Without Cancer (OHD-AD)	Concurrent Use of Opioids and Benzodiazepines (COB-AD)	Oral Evaluation During Pregnancy: Ages 21 to 44 (OEVP-AD)	Ambulatory Care Sensitive ED Visits for Non-Traumatic Dental Conditions in Adults (EDV-AD)	CAHPS Health Plan Survey 5.1H, Adult Version (Medicaid) (CPA-AD)	Long-Term Services and Supports Comprehensive Care Plan and Update (CPU-AD)	National Core Indicators Survey (NCIIDD-AD)	Postpartum Depression Screening and Follow-Up: Age 21 and Older (PDS-AD)**
Texas	34	1	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	X	-	-	X	-	X	X
U.S. Virgin Islands	1	0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	X	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Utah	32	0	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	X	X	X	X	X	-	X	X	X	X	X	X	X	X	-	-	X	X	-	X	-	X	-
Vermont	31	0	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	X	-	X	X	X	X	X	X	X	X	X	X	X	X	-	X	X	-	-	X	-	-	-
Virginia	14	0	X	X	X	X	X	X	X	X	X	X	X	-	-	-	-	-	-	-	X	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	X	-	X	-
Washington	34	0	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	-	X	-	X	-	X	-
West Virginia	33	0	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	X	X	X	X	X	X	X	X	X	X	X	X	-	X	X	-	X	X	-	-	-
Wisconsin	18	0	X	X	X	X	X	X	X	X	X	X	X	-	-	X	-	-	X	X	X	-	-	-	-	-	-	-	-	-	-	-	X	-	-	-	-	X	-	X	-
Wyoming	32	0	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	-	X	X	X	-	X	X	X	X	-	X	X	X	X	X	X	-	-	X	-	X	-

Source: Mathematica analysis of the Quality Measure Reporting (QMR) system reports for the 2025 Core Set as of April 28, 2026; Centers for Disease Control and Prevention Wide-ranging Online Data for Epidemiologic Research (CDC WONDER) data as of October 30, 2025; National Core Indicators (NCI) data submitted by states to the National Association of State Directors of Developmental Disabilities Services (NASDDDS) and the Human Services Research Institute (HSRI) (The NCI National Team) through the Online Data Entry System (ODESA) for the July 2024 to June 2025 data collection period, corresponding to the 2025 Adult Core Set, as of May 27, 2026; and Agency for Healthcare Research and Quality (AHRQ) and Centers for Medicare & Medicaid Services (CMS) analysis of the AHRQ CAHPS Database for the July 2024 to June 2025 data collection period as of January 20, 2026. CMS used two data sources for reporting the MSC-AD measure for the 2025 Adult Core Set. The two sources are the AHRQ CAHPS Database and QMR system reports for the 2025 Core Set.

Notes: The term “states” includes the 50 states, the District of Columbia, Guam, Puerto Rico, and the U.S. Virgin Islands.

CAHPS = Consumer Assessment of Healthcare Providers and Systems; ED = Emergency Department; HbA1c = Hemoglobin A1c

The 2025 Adult Core Set includes 38 measures. Information about the updates to the 2025 Core Sets is available at 2025 Updates to the Child and Adult Core Health Care Quality Measurement Sets and Mandatory Reporting Guidance. This table includes all 2025 Adult Core Set measures, including measures that were reported by states using “other specifications” and measures for which the rates are not publicly reported due to CMS data suppression rules.

*Starting with the 2024 Core Set, reporting of the behavioral health measures on the Adult Core Set is mandatory for the 50 states, the District of Columbia, Guam, Puerto Rico, and the U.S. Virgin Islands. Measures marked with an asterisk are mandatory for 2025 Adult Core Set reporting. More information is available in the Mandatory Reporting Final Rule and Initial Core Set Mandatory Reporting Guidance State Health Official Letter.

**The PDS-AD measure is a provisional measure for 2025 reporting. Provisional measures are voluntary and are not considered part of the 2025 Adult Core Set. For more information, see https://www.medicaid.gov/federal-policy-guidance/downloads/sho24001.pdf.

X = measure was reported by the state; - = measure was not reported by the state.